



Viewpoint

Quantification of tissue optical properties: perspectives for precise optical diagnostics, phototherapy and laser surgery

Alexey N Bashkatov

Research-Education Institute of Optics and Biophotonics, Saratov National Research State University, Saratov, Russia
Interdisciplinary Laboratory of Biophotonics, National Research Tomsk State University, Tomsk, Russia

E-mail:

a.n.bashkatov@mail.ru

Elina A Genina

Research-Education Institute of Optics and Biophotonics, Saratov National Research State University, Saratov, Russia
Interdisciplinary Laboratory of Biophotonics, National Research Tomsk State University, Tomsk, Russia

E-mail: eagenina@yandex.ru

Vyacheslav I Kochubey

Research-Education Institute of Optics and Biophotonics, Saratov National Research State University, Saratov, Russia
Interdisciplinary Laboratory of Biophotonics, National Research Tomsk State University, Tomsk, Russia

E-mail: saratov_gu@mail.ru

Valery V Tuchin^{1,2,3}

¹Research-Education Institute of Optics and Biophotonics, Saratov National Research State University, Saratov, Russia

²Interdisciplinary Laboratory of Biophotonics, National Research Tomsk State University, Tomsk, Russia

³Institute of Precision Mechanics and Control of the Russian Academy of Sciences, Saratov, Russia

E-mail: tuchinvv@mail.ru

This Viewpoint relates to an article by A N Bashkatov *et al* (2005 *J. Phys. D: Appl. Phys.* **38** 2543) and was published as part of a series of Viewpoints celebrating 50 of the most influential papers published in the *Journal of Physics* series, which is celebrating its 50th anniversary.

Recent technological advancements in photonics have encouraged remarkable progress towards the development of innovative methods and systems for clinical functional optical imaging, laser surgery, and phototherapy. The development of the optical biomedical methods and techniques has stimulated a great interest towards the investigation of optical properties of human tissues, which define the efficacy and the impact of tissue optical probing via the possibility to predict precisely the photon propagation trajectories and the fluence rate distribution within irradiated tissues.

Examples of the application of measured tissue optical properties for diagnostics include monitoring of blood oxygenation and tissue metabolism [1, 2], imaging of skin and mucous cancer tissues [3–5], tracking of drug [6, 7], micro-container [8–10] and nanoparticle delivery [11, 12], etc. In therapeutics, the methods of photodynamic therapy (PDT) [5, 13], low-level laser therapy (LLLT) [14–16], photo-thermal therapy (PTT) [17] or plasmonic photo-thermal therapy (PPTT) [18, 19], need a careful dosimetry that is impossible to provide without quantification of the tissue optical properties. The precision of laser tissue ablation [8, 9], coagulation [20] and welding [21, 22] strongly depends on spectral properties of tissues. Therefore, for all these applications, the knowledge of tissue optical properties is of great importance for the explanation and the quantification of the diagnostic data, and for the prediction of light distribution and absorbed energy for therapeutic and surgical use.

For our knowledge, the quantitative study of the optical properties of tissues began in the fifties of the last century. However, the most intensive researches were done from the nineties to the present. A rapid surge in the publication of studies during this period was caused by the development of optical technologies applicable for non-invasive or least-invasive diagnostics, therapy, and surgery of different diseases, as well as the perfection of the techniques for measurements of optical parameters of tissues in a wide wavelength range.

From these studies, several books, book chapters, and reviews summarizing and analyzing data on optical properties can be mentioned [23–30]. The analysis of these references (and literature within them) shows that the optical properties of human and animal skin, subcutaneous adipose tissue, breast tissue, myocardium, muscle, the skull, *dura mater* and other brain/head tissues, maxillary sinuses, the mucous membrane, stomach wall mucosa, the colon, the peritoneum, the uterus, gallbladder tissue, the liver, the aorta, lung tissue, the sclera, the conjunctiva, the retina and other ocular tissues, hairs, bone, cartilage, tooth enamel and dentin, and blood have been investigated in the visible and near-infrared spectral ranges.

Among papers, dealing with skin optics, the paper of Bashkatov *et al* [31] is one of the most cited. The paper has 450 citations in the database ‘Scopus’, 401 citations in the database the ‘Web of Science’, 637 citations in Google Scholar, and 422 citations in the Russian Science Citation Index. The citations show a variety of advanced biomedical technologies which are in direct need of knowledge of the optical parameters of tissues.

As the most impressive and most useful examples of tissue quantified data application, we can present the dosimetry of radiation during PDT and PTT [5, 13–17] and laser surgery

[8, 9, 20–22], different imaging technologies [1–5], the development of batteryless solar-powered cardiac pacemakers [32, 33], standardization of tissue-mimicking phantoms [34, 35], assessment of tissue chromophores concentration and distribution [36, 37], and proposition of the adequate optical models of tissues [38–41].

In the nearest future, we are expecting a further increase of the number of publications related to quantification of optical properties not only healthy, but also pathologically modified tissues and intensive use of these data for more precise diagnostics and monitoring of malignant neoplasm, diabetes, cardio-vascular diseases, as well as for light dosimetry at PDT, LLLT, PTT, PPTT, and laser surgery.

More efficient, accurate, and fast computer algorithms and experimental methods which are already available or will appear soon will allow for the measurement of the optical properties of tissues *in vivo* in a real time. This will bring the optical method to a new horizon and will significantly improve the care of people's health. For example, quantification of time varying optical parameters caused by endogenous metabolic processes or the external impact on a tissue via thermal action or due to immersion or compression optical clearing [42, 43] will be possible.

New 'tissue diagnostic/therapeutic windows' in the NIR attracted a lot of attention recently [44, 45]. In addition to visible/NIR 'windows' such as I (625–975 nm), II (1100–1350 nm), III (1600–1870 nm), and IV (2100–2300 nm) [12, 13], a narrow UV 'window' (350–400 nm) can be also used [14]. Due to great interest to UV [46], MIR [47] and terahertz [48] ranges in biophotonics, new studies of the optical properties for different tissues in these prospective wavelength ranges are expected to be done soon.

In addition, we should also expect the growth of the application of optical methods for characterization of tissues for the aims of the food industry to test meat quality and freshness, as this is evident from recent publications [49, 50].

Acknowledgments

The work was carried out under the financial support from the Russian Science Foundation, grant No. 14-15-00186.

References

- [1] Huang J *et al* 2015 Second derivative multispectral algorithm for quantitative assessment of cutaneous tissue oxygenation *J. Biomed. Opt.* **10** 036001
- [2] McEwen M P *et al* 2009 Vessel calibre and haemoglobin effects on pulse oximetry *Physiol. Meas.* **30** 869–83
- [3] Cugmas B *et al* 2015 Detection of canine skin and subcutaneous tumors by visible and near-infrared diffuse reflectance spectroscopy *J. Biomed. Opt.* **20** 037003
- [4] Lu G *et al* 2014 Spectral-spatial classification for noninvasive cancer detection using hyperspectral imaging *J. Biomed. Opt.* **19** 106004
- [5] Valentine R M *et al* 2011 Monte Carlo modeling of *in vivo* protoporphyrin IX fluorescence and singlet oxygen production during photodynamic therapy for patients presenting with superficial basal cell carcinomas *J. Biomed. Opt.* **16** 048002
- [6] Caduff A *et al* 2010 Cutaneous blood perfusion as a perturbing factor for noninvasive glucose monitoring *Diabetes Technol. Ther.* **12** 1–9
- [7] Darvin M E *et al* 2012 Comparison of two methods for noninvasive determination of carotenoids in human and animal skin: Raman spectroscopy versus reflection spectroscopy *J. Biophotonics* **5** 550–8
- [8] Belikov A V *et al* 2015 Multi-beam laser-induced hydrodynamic shock waves used for delivery of microparticles and liquids in skin *Lasers Surg. Med.* **47** 723–36
- [9] Genina E A *et al* 2016 Fractional laser microablation of skin: increasing the efficiency of transcutaneous delivery of particles *Quantum Electron.* **46** 502–9
- [10] Svenskaya Yu I *et al* 2016 Photodynamic therapy platform based on localized delivery of photosensitizer by vaterite submicron particles *Colloids Surf. B* **146** 171–9
- [11] Zhou L *et al* 2015 Effects of titanium dioxide nanoparticles coupled with diode laser on optical properties of *in vitro* normal and cancerous human lung tissues studied with optical coherence tomography and diffuse reflectance spectra *J. Biomed. Opt.* **20** 046003
- [12] SalmanOgli A and Rostami A 2012 Modeling and improvement of breast cancer site temperature profile by implantation of onion-like quantum-well heteronano-crystal in tumor site *IEEE Trans. Nanotechnol.* **11** 1183–91
- [13] Montazerabadi A R *et al* 2012 The effects of combined treatment with ionizing radiation and indocyanine green-mediated photodynamic therapy on breast cancer cells *J. Photochem. Photobiol. B* **109** 42–9
- [14] Kim S *et al* 2015 Effects of tissue water content on the propagation of laser light during low-level laser therapy *J. Biomed. Opt.* **20** 051027

- [15] Kim S and Jeong S 2014 Effects of temperature-dependent optical properties on the fluence rate and temperature of biological tissue during low-level laser therapy *Lasers Med. Sci.* **29** 637–44
- [16] Bhowmik A *et al* 2014 Analysis of radiative signals from normal and malignant human skins subjected to a short-pulse laser *Int. J. Heat Mass Transfer* **68** 278–94
- [17] Henderson T A and Morries L D 2015 Near-infrared photonic energy penetration: can infrared phototherapy effectively reach the human brain? *Neuropsychiatric Dis. Treat.* **11** 2191–208
- [18] Soni S *et al* 2013 Role of optical coefficients and healthy tissue-sparing characteristics in gold nanorod-assisted thermal therapy *Int. J. Hyperth.* **29** 87–97
- [19] Lalonde B S-L *et al* 2013 Visible and near infrared resonance plasmonic enhanced nanosecond laser optoporation of cancer cells *Biomed. Opt. Express* **4** 490–9
- [20] Schweinsberger G R *et al* 2011 Noninvasive laser coagulation of the human vas deferens: optical and thermal simulations *Lasers Surg. Med.* **43** 443–9
- [21] Tabakoglu H O and Gulsoy M 2010 *In vivo* comparison of near infrared lasers for skin welding *Lasers Med. Sci.* **25** 411–21
- [22] Hu L *et al* 2011 Closure of skin incisions by laser-welding with a combination of two near-infrared diode lasers: preliminary study for determination of optimal parameters *J. Biomed. Opt.* **16** 038001
- [23] Cheong W-F *et al* 1990 A review of the optical properties of biological tissue *IEEE J. Quantum Electron.* **26** 2166–85
- [24] Mobley J *et al* 2015 Optical properties of tissues *Biomedical Photonics Handbook* ed T Vo-Dinh (Boca Raton, FL: CRC Press) pp 23–122
- [25] Bashkatov A N *et al* 2011 Tissue optical properties *Handbook of Biomedical Optics* ed D A Boas *et al* (Boca Raton, FL: CRC Press) pp 67–100
- [26] Sandell J L *et al* 2011 A review of *in vivo* optical properties of human tissues and its impact on PDT *J. Biophotonics* **4** 773–87
- [27] Bashkatov A N *et al* 2011 Optical properties of skin, subcutaneous, and muscle tissues: a review *J. Innov. Opt. Health Sci.* **4** 9–38
- [28] Jacques S L 2013 Optical properties of biological tissues: a review *Phys. Med. Biol.* **58** R37–61
- [29] Bosschaart N *et al* 2014 A literature review and novel theoretical approach on the optical properties of whole blood *Lasers Med. Sci.* **29** 453–79
- [30] Tuchin V V 2015 *Tissue Optics: Light Scattering Methods and Instruments for Medical Diagnosis* 3rd edn (Bellingham: SPIE Press) PM 254
- [31] Bashkatov A N *et al* 2005 Optical properties of human skin, subcutaneous and mucous tissues in the wavelength range from 400 to 2000 nm *J. Phys. D: Appl. Phys.* **38** 2543–55
- [32] Haeblerlin A *et al* 2014 Successful pacing using a batteryless sunlight-powered pacemaker *Europace* **16** 1534–9
- [33] Haeblerlin A *et al* 2015 The first batteryless, solar-powered cardiac pacemaker *Heart Rhythm* **12** 1317–23
- [34] Lemailet P *et al* 2015 Double-integrating-sphere system at the National Institute of Standards and Technology in support of measurement standards for determination of optical properties of tissue-mimicking phantoms *J. Biomed. Opt.* **20** 121310
- [35] Wrobel M S *et al* 2015 Multi-layered tissue head phantoms for noninvasive optical diagnostics *J. Innov. Opt. Health Sci.* **8** 1541005
- [36] Tseng S-H *et al* 2009 Chromophore concentrations, absorption and scattering of human skin *in vivo* *Opt. Express* **17** 14599–617
- [37] Chen Y-W *et al* 2016 Toward reliable retrieval of functional information of papillary dermis using spatially resolved diffuse reflectance spectroscopy *Biomed. Opt. Express* **7** 542–58
- [38] Bhandari A *et al* 2011 Modeling optical properties of human skin using Mie theory for particles with different size distribution and refractive index *Opt. Express* **19** 14549–67
- [39] Schneiderheinze D H P *et al* 2007 Modified discrete particle model of optical scattering in skin tissue accounting for multiparticle scattering *Opt. Express* **15** 15002–10
- [40] Bjorgan A *et al* 2014 Estimation of skin optical parameters for real-time hyperspectral imaging applications *J. Biomed. Opt.* **19** 066003
- [41] Yudovsky D and Pilon L 2010 Rapid and accurate estimation of blood saturation, melanin content, and epidermis thickness from spectral diffuse reflectance *Appl. Opt.* **49** 1707–19
- [42] Zhu D *et al* 2013 Recent progress in tissue optical clearing *Laser Photonics Rev.* **7** 732–57
- [43] Genina E A *et al* 2015 Optical clearing of biological tissues: prospects of application in medical diagnostics and phototherapy (Review) *J. Biomed. Photonics Eng.* **1** 22–58
- [44] Smith A M *et al* 2009 Second window for *in vivo* imaging *Nat. Nanotechnol.* **4** 710–11
- [45] Shi L *et al* 2016 Transmission in near-infrared optical windows for deep brain imaging *J. Biophotonics* **9** 38–43
- [46] Yao D-K *et al* 2012 Optimal ultraviolet wavelength for *in vivo* photoacoustic imaging of cell nuclei *J. Biomed. Opt.* **17** 056004
- [47] Walsh M J *et al* 2012 Attenuated total reflectance Fourier-transform infrared spectroscopic imaging for breast histopathology *Vib. Spectrosc.* **60** 23–8
- [48] Nazarov M M *et al* 2008 Terahertz time-domain spectroscopy of biological tissues *Quantum Electron.* **38** 647–54
- [49] Zamora-Rojas E *et al* 2013 Prediction of fatty acids content in pig adipose tissue by near infrared spectroscopy: at-line versus *in situ* analysis *Meat Sci.* **95** 503–11
- [50] Lu R (ed) 2016 *Light Scattering Technology for Food Property, Quality and Safety Assessment* (Boca Raton, FL: CRC Press)

Nowadays, dynamically developing optical (photonic) technologies play an ever-increasing role in medicine. Their adequate and effective implementation in diagnostics, surgery, and therapy needs reliable data on optical properties of human tissues, including skin. This paper presents an overview of recent results on the measurements and control of tissue optical properties. The issues reported comprise a brief review of optical properties of biological tissues and efficacy of optical clearing (OC) method in application to monitoring of diabetic complications and visualization of blood vessels a ABSTRACT: The determination of the tissue optical parameters and the light propagation in multicomponent tissues is an important tool for real-time laser treatment dosimetry (selective laser ablation, PDT) and non-invasive laser diagnostics (laser induced fluorescence spectroscopy, Save to Library.Â ABSTRACT: The determination of the tissue optical parameters and the light propagation in multicomponent tissues is an important tool for real-time laser treatment dosimetry (selective laser ablation, PDT) and non-invasive laser more. ABSTRACT: The